

The precession of the giant HH34 outflow: a possible jet deceleration mechanism

E. Masciadri^{1,2}, E. M. de Gouveia Dal Pino³, A. C. Raga⁴, A. Noriega-Crespo⁵

ABSTRACT

The giant jets represent a fundamental trace of the historical evolution of the outflow activity over timescales of $\sim 10^4$ yr, i.e. a timescale comparable to the accretion time of the outflow sources in their main protostellar phase. The study of such huge jets provides the possibility of retrieving important elements related to the life of the outflow sources. In this paper, we study the role of precession (combined with jet velocity-variability and the resulting enhanced interaction with the surrounding environment) as a deceleration mechanism for giant jets using a numerical approach. This thesis was proposed for the first time by Devine et al. (1997) but it could not be numerically explored until now because it is intrinsically difficult to reproduce, at the same time, the large range of scales from ~ 100 AU up to a few parsecs. In the present paper, we obtain predictions of H α intensity maps and position-velocity diagrams from 3D simulations of the giant HH 34 jet (including an appropriate ejection velocity time-variability and a precession of the outflow axis), and we compare them with previously published observations of this object. Our simulations represent a step forward from previous numerical studies of HH objects, in that the use of a 7-level, binary adaptive grid has allowed us to compute models which appropriately cover all relevant scales of a giant jet, from the ~ 100 AU jet radius close to the source to the ~ 1 pc length of the outflow. A good

¹Instituto de Astronomía, UNAM, Apdo. Postal 70-264, 04510 México, D. F., México, email: elena@astroscu.unam.mx

²Max-Planck Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany, email: masciadri@mpia.de

³Instituto Astronomico e Geofísico (IAG-USP), Universidade de São Paulo, Cidade Universitaria, R. do Matao, 1226, São Paulo, SP 05508-090, Brasil, email: dalpino@astro.iag.usp.br

⁴Instituto de Ciencias Nucleares, UNAM, Apdo. Postal 70-543, 04510 México, D. F., México, email: raga@astroscu.unam.mx

⁵SIRTF Science Center, California Institute of Technology, IPAC 100-22, Pasadena, CA 91125, email: alberto@ipac.caltech.edu

qualitative and quantitative agreement is found between the model predictions and the observations, indicating that a precession of the jet axis can indeed be the probable cause of the deceleration of the giant jets. Moreover, we show that a critical parameter for obtaining a better or worse agreement with the observations is the ratio ρ_j/ρ_a between the jet and the environmental densities. The implications of this result in the context of the current star formation models are discussed

1. Introduction

Herbig-Haro (HH) objects are the optical manifestations of outflows from young stellar objects (YSOs). Following the discovery of jet-like structures in HH objects (Dopita et al. 1982; Mundt & Fried 1983), many of these HH jets were observed in the Orion (Reipurth et al. 1986; Mundt et al. 1987; Reipurth 1989a, Reipurth 1989b) and Taurus (Mundt et al. 1988) star formation regions. These objects present a characteristic morphology of aligned knots extending over ~ 0.3 pc. It appears that the kinematics and morphologies of these jets depend simultaneously on the time-dependent nature of the outflow activity and on the interaction of the hypersonic flows with the surrounding interstellar medium.

It has recently been discovered (Bally & Devine 1994; Reipurth et al. 1997; Devine et al. 1997) that a few HH jets extend over distances of a few parsecs. For example, HH 111 shows a total extent of ≈ 7.7 pc, HH 34 of ≈ 3 pc and HH 355 a total extent of ≈ 1.55 pc.

Apart from their alignments, the main evidence that the knots belong to the same jet (and not to other, smaller outflows) is their kinematic association with red- and blue-shifted bipolar lobes. From the radial velocity, the proper motions and the distance of the knots from the source it has been possible to estimate a typical dynamical age of $\sim 10^4$ yr for these “giant jets”.

An important characteristic of the giant jets is that they appear to slow down for increasing distances from the outflow source. This effect is seen in the HH 34 (Devine et al. 1997) and in the HH 111 giant jets (Reipurth et al. 1997; Rosado et al. 1999). The present paper is concerned with the possible theoretical interpretations of the deceleration effect. This is a critical point in the determination of the jet’s age and also in the identification of the physical properties of the central engine that feeds the outflows.

The potential causes that might produce such a deceleration can be divided into two categories :

- internal causes, i.e. that the mechanism is intrinsically related to the properties of the outflow, for example to a temporal variability of the ejection velocity,
- external causes, i.e. that the deceleration is due to the drag effect resulting from the interaction of the hypersonic flow with the interstellar medium.

Previous studies have considered both of these possibilities. Cabrit & Raga (2000) considered the case of an ejection velocity which monotonically grows as a function of time, and tried to fit the observed, position-dependent jet velocity with different, parameterized forms of the ejection velocity time-dependence. These authors concluded that the only way to fit the observed kinematics of the HH 34 giant jet is with an ejection velocity that slowly increases over $\sim 5 \times 10^4$ yr, followed by a very strongly increasing ejection velocity over the last $\sim 10^4$ yr.

Cabrit & Raga (2000) argued that this very dramatic increase in the ejection velocity at recent times appeared to be unlikely, and then studied an alternative scenario. Following the idea proposed by Devine et al. (1997), they considered the knots along the HH 34 giant jet as scattered “bullets” resulting from the combination of an ejection velocity variability and a precession of the jet axis. Raga & Biro (1993) have carried out a theoretical study of this kind of flow, obtaining an analytic description of this “machine gun jet” flow and comparing this model with a numerical simulation of a radiative, 2D “slab” jet with a time-dependent ejection velocity and direction.

One could argue that the jet/counterjet symmetry observed in the HH 34 giant jet (see Devine et al. 1997) goes against the “environmental drag” scenario for the deceleration of this object. As there is no reason to suppose that the environments within which the jet and the counterjet are traveling have identical densities, one would think that the drag would introduce asymmetries between the two outflow lobes. However, the deceleration induced by the environmental drag is proportional to $\rho_a^{1/3}$ (where ρ_a is the environmental density, see Cabrit & Raga 2000), so that these asymmetries might not be so important.

Also, de Gouveia Dal Pino (2001) presented 3D simulations (done with the Lagrangian SPH method) of HH 34 assuming a sinusoidal ejection velocity variability of the kind used by Raga & Noriega-Crespo (1998) and studying the two cases of a pressured and overpressured jet without considering the precession contribution. To reproduce the large spatial working surface structures the author used a half-amplitude of the velocity modulation of $\approx 100 \text{ km s}^{-1}$ and a period of 760 yr. The author obtained encouraging results showing that a deceleration of the jet velocity was obtained from the model (lower decelerations being produced for initially overpressured jets than for pressure matched jets). She concluded from the simulations that the deceleration was related to the temporal

velocity variability of the jet at injection and was mainly caused by progressive momentum transfer sideways into the surrounding medium by the expelled gas from the travelling working surfaces. She also found that a steady state jet, with similar initial conditions to those of the pulsed jet, experienced, on the contrary, an initial acceleration followed by a constant velocity propagation regime, which was an additional indication that the primary source of deceleration in giant flows could not be attributed to simple breaking of the jet head against the external medium. These models were run over a distance of only 0.3 pc, well below the size of the giant HH 34 jet (see Devine et al. 1997).

In the present paper, we carry out a numerical study of the effect of a precession of the outflow axis on the deceleration of a giant jet. The existence of such a phenomenon is suggested by the observed morphology of the flow, which shows evidence of long period (of the order of 10^4 yr) precession (Devine et al. 1997). The precession is generally ascribed to tidal forces produced by a companion in a binary or multiple system. Even though in the case of HH 34 the binary source has not yet been resolved, there are elements that indicate that this source could be a binary such as the discovery of a second outflow (HH 534) emanating from the source as well as the abrupt change of direction of the jet axis near the knots B (Reipurth et al. 2002). Less evident is the explanation of the existence of a precession period of the order of 10^4 yr.

As reported in Terquem et al. (1999), the precession period (τ_p) depends on the orbital parameters and the spatial extent of the accretion disk. The precession period is generally at least one order of magnitude larger than the orbital period. Therefore, to justify such a large τ_p we need a value of the ratio between the accretion radius disk R and the orbital radius r_0 ($\sigma = R/r_0$) of the order of 10^{-3} (see Masciadri & Raga, 2002).

Reipurth (2000) has also proposed that perturbations on an accretion disk due to close passages (i. e., at perihelion) of a binary companion in an elliptical orbit could be responsible for producing a time-variability in the ejection of the outflow. It is of course unclear whether or not such a mechanism could produce a variability in the ejection velocity such as the one included in our jet models.

We carry out the simulations of only one of the two lobes of HH 34 (with and without precession) conserving the same geometrical and physical parameters. However, our numerical simulations could correspond to any of the two lobes of the HH 34 giant jet. Actually, our simulations are made to reproduce the morphology of the northern lobe of the HH 34 outflow, and in order to compare the predicted maps with the southern lobe it is necessary to carry out a point reflection of the predicted maps with respect to the position of the outflow source. This point symmetry of a precessing jet/counterjet system is clearly seen in the observations of the HH 34 giant outflow (Devine et al. 1997).

We underline that this study cannot be applied in a simple way to other giant jets. The two lobes of HH 111, for example, show quite straight paths. HH 355, on the contrary, seems to precess with a half-opening angle of $\sim 13^\circ$ and a period of ~ 1500 - 2000 yr (Reipurth et al. 2002). Further work should be carried out to study the properties of these other giant flows.

We find that the impact of the precession on the deceleration mechanism is quite considerable, and that the dynamical age of the jet grows by ~ 3000 yr when including a precession. We also find that H α intensity maps and position-velocity diagrams obtained from models with precession reproduce the observations of the HH 34 giant jet in a qualitatively successful way (an agreement which is not found for models without precession).

We add that, from a numerical point of view it is not a simple exercise to reproduce the evolution of outflows over such a large spatial and temporal extent particularly given the small initial radius of the beam. Indeed, in order to cover the whole domain of 1.5 pc, previously published simulations used $r_j = 10^{16}$ cm (de Gouveia Dal Pino 2001) and $r_j = 10^{17}$ cm jet radii (Cabrit & Raga 2000), which are 1-2 order of magnitude greater than the width of the jet as observed in HST images (Reipurth & Raga 1999). One of the goals of our study is to simulate the HH 34 giant jet over its full extent (~ 1.5 pc) using the correct $r_j = 3 \times 10^{15}$ cm initial jet radius corresponding to $0''.4$ at 460 pc. We underline that the radius is measured at $\sim 10''$ distance away from the HH 34 source.

In Section 2, we describe the parameters and the ejection velocity time-variability used in our models of the HH 34 jet. In Section 3 the numerical simulations are discussed, and H α maps and position-velocity diagrams predicted from models with and without precession are presented and compared with the corresponding observations of the HH 34 giant jet. In Section 4 we summarize the conclusions of this study.

2. Parameters for the jet models

Following the study of HH 34 of Raga & Noriega-Crespo (1998), we attempt to reproduce the structure of this jet by assuming the existence of a sinusoidal ejection velocity time-variability. As our work is focussed on trying to reproduce the large scale structure of this outflow, a single mode time-variability is appropriate, as opposed to the three-mode variability used by Raga & Noriega-Crespo (1998) to model the structures close to the source along the southern lobe.

We therefore consider an ejection velocity of the form

$$v(t) = v_j + v_a \sin \omega_a t \quad (1)$$

where v_j is the average jet velocity, $\omega_a = 2\pi/\tau_a$ is the frequency (τ_a is the period) and v_a is the half-amplitude of the variability law.

We adopt the following values : $v_j = 300 \text{ km s}^{-1}$, $v_a = 110 \text{ km s}^{-1}$ and $\tau_a = 1010 \text{ yr}$. These values are consistent with the long period mode deduced from the kinematics of the southern lobe of HH 34 by Raga & Noriega-Crespo (1998). Also, these parameters give a spatial separation

$$\Delta x \approx \tau_a v_j = 9.5 \times 10^{17} \text{ yr}, \quad (2)$$

which corresponds to a separation of $\approx 120''$ on the plane of the sky (considering a distance of 460 pc to HH 34 and a $\phi = 28^\circ$ angle between the outflow axis and the plane of the sky). This separation is consistent with the separations between successive knots along the HH 34 giant jet (see Devine et al. 1997). Also, the images of the HH 34 giant jet show evidence of a long period precession. From the images of Bally & Devine (1994) and Devine et al. (1997), we find that both the jet and the counterjet appear to lie within a cone of half-opening angle $\alpha = 6^\circ$, and that the locus of the jet appears to imply a $\tau_p = 12000 \text{ yr}$ precession period. We take these values as estimates of a possible precession of the HH 34 outflow.

For our numerical simulations, we consider that the jet has a top-hat initial cross section, of radius $r_j = 3 \times 10^{15} \text{ cm}$, corresponding to $0''.4$ at the distance of HH 34, which is consistent with the recent results obtained from HST images of this object (Reipurth et al. 2002). Moreover, we choose two initial jet number densities, $n_j = 10^3$ and $5 \times 10^3 \text{ cm}^{-3}$. We have assumed that the surrounding environment is homogeneous, with a $n_a = 100 \text{ cm}^{-3}$ number density. We have set the initial temperature of both the jet and the environment to 1000 K, and assumed that the gas is neutral, with the exception of Carbon, which is singly ionized.

3. Numerical simulations

3.1. The numerical computations

The simulations were carried out using 3D gasdynamic adaptive grid yguazu-á code. This code integrates the 3D gasdynamic equations and a set of atomic/ionic reaction equations for the species HI, HII, OI, OII, OIII, CII, CIII and CIV. They are the main contributors to the cooling function. The yguazu-á code employs the *flux-vector splitting*

algorithm of Van Leer (1982) and it was described in detail by Raga et al. (2000). The reaction rates and the non-equilibrium cooling function that we have used are given by Raga et al. (2002a).

The computations were carried out on a 7-level, binary adaptive grid with a maximum resolution along the three axes of 1.95×10^{15} cm. The computational domain extends over $10^{18} \times 10^{18} \times (4 \times 10^{18})$ cm, corresponding to $512 \times 512 \times 2048$ grid points at the highest resolution grid level. The jet is injected at the origin in the centre of the xy -plane and the outflow axis precesses around the z -axis.

The maximum resolution (level 7) is allowed only within a region limited within a spherical surface of radius 0.5×10^{18} cm centred on the injection point. The next highest resolution (level 6) is allowed only in a region with an outer radius of 1.5×10^{18} cm.

Figure 1 shows the adaptive grid configuration on the xz -plane obtained from Model B (see below) after a $t = 8.4 \times 10^3$ s time integration. This figure shows the grid in a domain of size $(10^{18}, 3 \times 10^{18})$ cm (left frame), and two successive zooms (centre and right) showing the higher resolution grids.

We computed three different models :

- **Model A** - a jet that precesses in a cone of half-angle $\alpha = 6^\circ$ with a period $\tau_p = 12000$ yr. The initial jet density has a $n_j = 10^3 \text{ cm}^{-3}$ value, resulting in a $\rho_j/\rho_a = 10$ jet to environment density ratio.
- **Model B** - the same precession as Model A, but a jet with $n_j = 5 \times 10^3 \text{ cm}^{-3}$ ($\rho_j/\rho_a = 50$),
- **Model C** - a jet with the same parameters as Model A but without precession.

The three models have the time-dependent ejection velocity given by Eq. (1), and the jet and ambient medium parameters given at the end of §2.

As an example of the flows that result from our simulations, in Figure 2 we show a time-sequence of the column density obtained for Model B (which, as we show below is the model that more closely resembles de HH 34 giant jet). The column density was obtained by integrating the density field along the y -axis. Figure 2 shows that many working surfaces are formed, and that they travel in different directions away from the source as more or less independent “bullets”. The qualitative features of this kind of flow were discussed by Raga & Biro (1993).

3.2. H α maps

All the three models (A, B and C) were run over different time intervals such that the outflows have traveled a distance of $\approx 3 \times 10^{18}$ cm, equivalent to ≈ 1 pc. Figures 3, 4 and 5 show the temporal evolution of the H α maps predicted from the three models. These maps were computed assuming an $\phi = 28^\circ$ angle between the outflow axis and the plane of the sky. The H α emission coefficient was calculated considering the contributions of the recombination cascade and of collisional excitations from the ground state.

It is evident that the loci of the jets are quite different for Models A, B and C. Model A and B show a bending not present in Model C. Moreover the first two models have a larger concentration of knots than the Model C. The non-precessing model C, of course, shows a structure of aligned knots that looks dramatically different from Models A and B. Also, the times t_A , t_B and t_C (corresponding to Models A, B and C, respectively) at which the jet heads reach a distance of 1 pc from the source are substantially different from each other. We obtain $t_A = 18.4 \times 10^3$ yr, $t_B = 8.4 \times 10^3$ yr and $t_C = 5.4 \times 10^3$ yr. We note that the maximum H α map value is equal to 7×10^{-6} erg cm $^{-2}$ s $^{-1}$ sr $^{-1}$ in Model A, equal to 2.5×10^{-5} erg cm $^{-2}$ s $^{-1}$ sr $^{-1}$ in Model B and equal to 2×10^{-7} erg cm $^{-2}$ s $^{-1}$ sr $^{-1}$ in Model C. Comparing the results of our simulations with the measured H α emission (Reipurth et al. 2002, Fig. 4) we conclude that the model that better reproduces the observed H α fluxes is Model B. Indeed, the highest H α contour obtained displayed in the HST images of Reipurth et al. (2002, Fig. 4) corresponds to $\approx [13-18] \times 10^{-18}$ erg cm $^{-2}$ s $^{-1}$ pixel $^{-1}$, which is equivalent to $[3-5] \times 10^{-4}$ erg cm $^{-2}$ s $^{-1}$ sr $^{-1}$. We conclude that the model that better reproduces the observed H α fluxes is Model B.

However, we find that the model predictions give H α maps that are in general one order of magnitude fainter than the the observed ones. In a previous study, Raga & Noriega-Crespo (1998) found that for an initial jet density of 5×10^2 cm $^{-3}$, the predicted flux is two orders of magnitude fainter than HH 34. The fact that we obtain a one order of magnitude higher flux is consistent with the fact that in the present model we have considered a 5×10^3 cm $^{-3}$ jet density (in other words, the models follow the standard intensity $\propto$ pre-shock density scaling law). Therefore, in order to obtain a better agreement with the H α fluxes of HH 34, we would need a model in which both the jet and the environment are denser by a factor of ~ 10 . We have not computed denser models, because they have shorter cooling distances, which would not be resolved appropriately in our numerical simulations.

At the same time, the morphology of the knot distribution in Model B better resembles the observed morphology of the HH 34 giant jet (Devine et al. 1997, Fig. 6) than do the knot structures predicted from Models A and C. We note that Model A (with

$\rho_j/\rho_a = 10$) produces, at a time $t \approx 11.2 \times 10^3$ yr from the beginning of the simulation, an extended shock structure (resulting from a piling up of several working surfaces) at about $z \approx 0.7 \times 10^{18}$ cm. Such piling ups of working surfaces do not occur in Model B (which has $\rho_j/\rho_a = 50$), resulting in quite dramatically different H α maps being predicted from both models. We therefore conclude that the jet-to-environment density ratio ρ_j/ρ_a is a critical parameter for jets from precessing sources, and has to be adjusted in order to be able to reproduce the observations of a given HH jet.

The calculation of the temporal evolution of the H α maps permitted us to retrieve the proper motions of the knots dispersed along the jet path. Figure 6 shows H α maps obtained at two different times. From the positions of the knots in these two frames, we have computed proper motion vectors (shown on the left plot). One can observe that the magnitude of the proper motion velocity slows down as a function of distance from the source. This result is qualitatively consistent with the proper motion measurements of Devine et al. (1997), which show tangential velocities which decrease from 198 *km/sec* to 96 *km/sec* as a function of distance from the source along the giant jet. We do not attempt to carry out a quantitative comparison of the predicted and observed proper motions, as the errors of the proper motions of the HH 34 giant jet are quite large.

In order to carry out a more quantitative comparison between the kinematics of HH 34 and our models, it is better to compare the predicted and measured radial velocities. This kind of comparison between models and observations is described in the following section.

3.3. Position-velocity diagrams and radial velocities

One of the most reliable ways to validate the simulations is to compare the PV diagrams obtained from the three models with observations. Indeed, one of the most interesting results of Devine et al. (1997, Fig. 7) are their quite accurate measurements of decreasing radial velocities as a function of distance from the source along the HH 34 giant jet. Figure 7 shows the PV diagrams predicted obtained from Models A, B and C. The black, thin line represents the maximum of the emission (as determined from quadratic fits to the line peaks) vs. distance from the source. The three PV diagrams are calculated at the times at which the head of the jet reaches 3×10^{18} cm (these times are t_A , t_B and t_C , see §3.2). The same $\phi = 28^\circ$ angle between the outflow axis and the plane of the sky was considered.

Figure 8 shows the radial velocity vs. distance from the source measured by Devine et al. (1997) for the knots along both the north and south lobes of the HH 34 giant jet. We

changed the sign of the velocities of the knots in the south lobe, so that all of the velocities are then positive. The bold points are the peak of the gaussian fit and the error bars represent the FWHM of the emission line. We omitted the point related to the HH 34 X knot because, as one can see in Figure 7 of Devine et al. (1997), it is quite distant from the trend defined by the velocities of all of the other knots in the outflow.

The three continuous lines join the radial velocities of the working surfaces (the sharp velocity jumps in Fig. 7) obtained from Model A (dotted line), Model B (thin line) and Model C (dot-dash-dot line). These radial velocities (shown with stars in the PV diagrams of Fig. 7) correspond to the successive knots which are seen in the predicted $H\alpha$ maps. The positions of the stars correspond to the intensity maxima (i. e., the knots) in the corresponding $H\alpha$ maps.

Both Figures 7 and 8 show that Model A produces too steep radial velocity vs. distance decrease, particularly beyond 2×10^{18} cm from the source. Model C produces radial velocities which are too high over the whole path of the jet. On the other hand, Model B produces radial velocities which agree quite well with the observed values.

Figure 8 (right-hand side) shows linear fits to the radial velocities as a function distance from the source obtained from the observations and from Models A, B and C, in other words, the lines trace the decreasing rate of the radial velocity. In this graph, we again see that while Model A produces a too sharp drop in radial velocities, and Model C gives velocities which are too high, Model B does produce a good agreement with the observations.

3.4. Effects of the precession vs. the initial jet to environmental pressure ratio

What can we say about the impact of the precession and of the initial jet to environmental pressure ratio ? We define η as the ratio between the initial jet and environmental density ($\eta = \rho_j / \rho_a$) and we define k as the corresponding pressure ratio ($k = p_j / p_a$). Knowing the values of ρ_a , ρ_j , T_a and T_j we can calculate $k = 10$ for the Models A and C and $k = 50$ for Model B. All three models are therefore overpressured, Model A and C have the same value of k and Model B is more highly overpressured. Figure 8 shows that the deceleration rates of Models A and C are very different in spite of the fact that the two models have identical k . Considering that in Model A we have a precession that is absent in the Model C we would then conclude that the difference in the deceleration is a direct result of the precession.

de Gouveia Dal Pino (2001) studied the effects of different k values on a non-precessing

jet and found that lower values of k resulted in stronger decelerations for the working surfaces. This is a result of the fact that the different k values are obtained by changing the environmental temperature T_a , with higher values of T_a resulting in a stronger coupling between the aligned working surfaces and the surrounding gas (and therefore causing a stronger deceleration). This rather subtle effect is not likely to be important in the case of a precessing jet for which the scattered working surfaces ram directly into the undisturbed environment. For this case, the important parameter for determining the deceleration rate is the density ratio η .

It would be interesting to study in the future the effect on the deceleration mechanism of two other flow parameters: the angle and the period of the precession. However, an idea of the effects that would be introduced can be obtained from the analytic model of Raga & Biro (1993). These authors showed that the distance (measured along the precession axis) at which the working surfaces become independent “bullets” interacting with the surrounding environment is $z_b \approx r_j \tau_p \cot \alpha / (\pi \tau_a)$, corresponding to $\approx 4 \times 10^{17}$ cm for our model B. Therefore, in our model, the $z < z_b$ region in which the working surfaces are sheltered from a direct interaction with the environment is small, so that they are subjected to the full environmental drag during most of their evolution. We would then expect that all models with α , τ_p and τ_a such that z_b is much smaller than the full length of the jet will have deceleration properties similar to our models, provided that they have the same r_j and jet-to-environment density ratios.

4. Conclusion

In this paper, we explore the role of a precession of the outflow axis on the deceleration of giant HH jets. In particular, we try to simulate the HH 34 giant jet, and we compare the results of our numerical simulations with previously published observations of this object.

Our simulations represent a step forward from previous numerical studies of HH objects, in that the use of a 7-level, binary adaptive grid has allowed us to compute models which appropriately cover all of the relevant scales of a giant jet, from the ~ 100 AU jet radius close to the source to the ~ 1 pc length of the outflow. Previous simulations of giant jets either did not cover the length of a real flow (de Gouveia dal Pino 2001), or else had a very large jet radius (Cabrit & Raga 2000 and de Gouveia dal Pino 2001).

A set of simulations done with and without precession of the outflow axis are presented, and predictions of $H\alpha$ maps, proper motions and radial velocities are compared with the observations of Devine et al. (1997). The principal conclusions of our study are the

following :

- we see that the morphology and kinematics of the HH 34 giant jet can be reproduced with a model of a jet with a sinusoidal ejection velocity variability (with a mean velocity of 300 km s^{-1} , a half-amplitude of 100 km s^{-1} and a period of 1010 yr) and a precession of the outflow axis (with a half-angle of 6° and a 12000 yr period). The simulated and measured $\text{H}\alpha$ maps and radial velocities show a good qualitative as well as quantitative agreement,
- comparing simulations done with and without precession we showed that the simple precession can give differences in the jet age estimations of the order of about 3000 yr. This proves that the drag effect produced by the external medium on the working surfaces is not negligible with respect to the $\sim 10^4$ yr dynamical timescale of the outflow.
- we proved that the ρ_j/ρ_a ratio is a critical parameter in the determination of the deceleration rate of the jet, and that it has to be properly adjusted in order to be able to fit the observed properties of a giant HH flow.

The results of our study do not exclude that, in other giant jets, a correct jet deceleration could be attained without the precession. More models should be tested with different ρ_j/ρ_a ratios to obtain more definite conclusions. Besides this, in the case of HH 34, the results seem to indicate that the precession has a fundamental role in the deceleration mechanism. We underline that the non-precessing, velocity-variable jet of Model C also decelerates, however it has radial velocities which are larger than the observed ones in HH34 (Devine et al. 1997). Previous 3D modeling of time-variable, non-precessing giant outflows (de Gouveia Dal Pino 2001) had also detected jet deceleration that reproduced the observations only qualitatively. Therefore, the results of the present work indicate that in the case of HH34, it is the combined effect of both, the jet temporal velocity variability and the precession (along with the appropriate choice of the ratio ρ_j/ρ_a) that reproduces the observed deceleration pattern in that source (as in Model B). One could argue that Model C with different ρ_j/ρ_a ratios or velocity variability law could reproduce the correct deceleration. On the other hand we observe that a change of the ρ_j/ρ_a ratio seems to produce a modification in the deceleration rate (Fig. 8 right hand side). A different velocity variability would produce a different distribution of the working surfaces along the jet trajectory and an $\text{H}\alpha$ map characterized by a different emission. We also recall that the emission of the $\text{H}\alpha$ maps strongly depends on the ρ_j/ρ_a ratio. A smaller ρ_j/ρ_a ratio in Model C would probably reproduce a $\text{H}\alpha$ map characterized by an emission level which is too low (see Section 3.2). It therefore appears that these high radial velocities are due

neither to an incorrect ρ_j/ρ_a ratio nor to a velocity variability law different from the one that we have considered.

In our simulations of the HH 34 giant jet, all of the structure of the outflow is due to a velocity variability with a single, sinusoidal mode and a well ordered precession. In our model, both the velocity variability and the precession last for all of the life of the outflow. Furthermore, the ejection velocity variability that we have used agrees with the one determined by Raga & Noriega-Crespo (1998) for the region between the source and HH 34S, so that there is evidence that this variability is continuing to at least quite close to the present time.

The existence of a precession is an indication that the source belongs to a binary or multiple system. This is in agreement with Reipurth (2000), who argued that the sources of giant HH jets are binary or multiple systems.

However, our results seem to be less consistent with the thesis proposed by Reipurth (2000) that giant jets are fossil records of the evolution of orbital motions in disintegrating multiple systems. This process has three distinct phases : (1) a non-hierarchical state (called *interplay*) in which the multiple system performs a random motion, (2) a close triple approach in which a close binary is formed and a low mass star or embryo moves over to a larger orbit and (3) an ejection phase in which the low mass embryo is ejected from the nucleus of the system.

In our models, the structure of the HH 34 giant jet is reproduced without the need of having different properties of the ejection at different times, reflecting qualitative changes in the outflow source resulting from the three phases of a disintegrating multiple system (see above). Therefore, we conclude that if the HH 34 system does correspond to an outflow history with distinct phases, the evidence for this appears to have been lost in the complexities of the interaction between the jet and the surrounding environment.

The work of EM and AR was supported by CONACyT grants 34566-E and 36572-E. We thank Israel Díaz for installing the new computer with which the numerical simulations were carried out. E.M.G.D.P. has been partially supported by grants of the Brazilian Agencies FAPESP (grant 97/13084-3) and CNPq. A.N-C. research was carried at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA; and partially funded by ADP grant NRA0001-ADP-096. We acknowledge an anonymous referee for giving us helpful comments.

REFERENCES

- Bally & Devine, 1994, ApJ, 428, L65
- Cabrit S. & Raga A., 2000, A&A, 354, 667
- de Gouveia Dal Pino E., 2001, ApJ, 551, 347
- De Young D.S. & Axford W.I., 1967, Nat., 216, 129
- Devine D., Bally J., Reipurth B., Heatcote S., 1997, AJ, 114, 2095
- Dopita M. A., Evans I., Schwartz R. D., 1982, ApJ, 263, L73
- Mundt R. & Fried, 1983, ApJ, 274, L83
- Mundt R., Brugel E.W., Buehrke T., 1987, ApJ, 319, 275
- Mundt R., Buehrke T., Ray T., 1988, ApJ, 333, L69
- Raga A., Cantó J., Binette L., Calvet N., 1990, ApJ, 364, 601
- Raga A. C., Biro S., 1993, MNRAS, 264, 758
- Raga A., Noriega-Crespo A., 1998, AJ, 116, 2943
- Raga A. C., Mellema G., Arthur S. J., Binette L., Ferruit P., Steffen W., 1999, RMxAA, 35, 123
- Raga A., Navarro-González R., Villagrán-Muniz M., 2000, RMxAA, 36, 67
- Raga A. C., de Gouveia dal Pino E. M., Noriega-Crespo A., Veázquez P. F., Mininni P., 2002a, A&A, submitted.
- Reipurth B., Bally J., Graham J.A., Lane A.P., Zealey W.J., 1986, A&A, 164, 51
- Reipurth B., 1989a, Nature, 340, 42
- Reipurth B., 1989b, A&A, 220, 249
- Reipurth B., Bally J., Devine D., 1997, AJ, 114, 2708
- Reipurth B. & Raga A., 1999, *Origin of Stars and Protoplanetary systems*, ed. C. Lada & N. D. Kylafis, NATO Sci. Ser. C, 540, 267
- Reipurth B., 2000, AJ, 120, 3177

- Reipurth B., Bally J., 2001, ARA&A, 39, 403
- Reipurth B., Heathcote S., Morse J., Hartigan P., Bally J., 2002, AJ, 123, 362
- Rosado M., Raga A. C., Arias L., 1999, AJ, 117, 462
- Terquem, C., Eislöffel, J., Papaloizou, J. C. B., Nelson, R. P., 1999, ApJ, 512, L131.
- Van Leer, ICASE Report, No.82-30, 1982,

Fig. 1.— 7 level, binary adaptive grid obtained from Model B after a 8.4×10^3 yr time integration. The abscissas and ordinates are labeled in units of 10^{18} cm. On the left hand side, the grid extended over the $(10^{18}, 3 \times 10^{18})$ cm domain is shown. In the centre and right hand side, successive zooms of smaller regions are shown. The points represent the central positions of the grid cells on the xz-plane.

Fig. 2.— Temporal evolution of the column density (integrated along the y -axis) obtained from Model B. The abscissas and ordinate are labeled in units of 10^{18} cm. The column densities are shown with the linear grey-scale given (in cm^{-2}) by the bar on the right.

Fig. 3.— Temporal evolution of the $\text{H}\alpha$ maps obtained from **Model A**. The last frame corresponds to the time at which the jet extends out to $\approx 3 \times 10^{18}$ cm from the source. The abscissas and ordinates are labeled in units of 10^{18} cm. The $\text{H}\alpha$ maps are shown with the logarithmic grey-scale given (in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$) by the bar on the right. The maximum $\text{H}\alpha$ intensity value is of $7 \times 10^{-6} \text{ erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$.

Fig. 4.— Temporal evolution of the $\text{H}\alpha$ maps obtained from **Model B**. The last frame corresponds to the time at which the jet extends out to $\approx 3 \times 10^{18}$ cm from the source. The abscissas and ordinates are labeled in units of 10^{18} cm. The $\text{H}\alpha$ maps are shown with the logarithmic grey-scale given (in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$) by the bar on the right. The maximum $\text{H}\alpha$ intensity is of $2.5 \times 10^{-5} \text{ erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$.

Fig. 5.— Temporal evolution of the $\text{H}\alpha$ maps obtained from **Model C**. The last frame corresponds to the time at which the jet extends out to $\approx 3 \times 10^{18}$ cm from the source. The abscissas and ordinates are labeled in units of 10^{18} cm. The $\text{H}\alpha$ maps are shown with the logarithmic grey-scale given (in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$) by the bar on the right. The maximum $\text{H}\alpha$ intensity is of $2 \times 10^{-7} \text{ erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$.

Fig. 6.— The left plot shows the $\text{H}\alpha$ map at 6×10^3 yr. The right hand side plot shows the $\text{H}\alpha$ map at 6.4×10^3 yr. On the left plot the proper motions of the four brightest knots are marked. The abscissas and ordinates are labeled in units of 10^{18} cm. The $\text{H}\alpha$ maps are shown with the logarithmic grey-scale given (in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$) by the bar on the left.

Fig. 7.— Position velocity diagram obtained for the Models A (left hand side), B (centre) and C (right hand side). The emission is shown with a logarithmic greyscale given (in $\text{erg s}^{-1} \text{cm}^{-1} (\text{cm/s})^{-1} \text{sr}^{-1}$). The black, thin line represents the emission maximum of the PV diagram vs. distance from the source. On the abscissas, the radial velocity values are given in 100 km/s units. The ordinates give the position along the spectrograph slit in 10^{18} cm units. The abscissas give the absolute values of the radial velocities.

Fig. 8.— On the left hand side are shown the absolute values of the radial velocity vs. distance from the source measured by Devine et al. (1997) (bold points and errors bar). The graph also shows the radial velocities obtained from Model A (dotted line), Model B (thin line) and Model C (dot-dash-dot line). On the right hand side, linear fits to the values reported in the left plot are shown.

This figure "f1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0208277v1>